


Review Article

Open Access

Millets: For Food and Nutritional Security**Tamanna Sharma¹, Nikhil Thakur², Jyoti Sharma³ and Monika Kumari⁴**¹Division of Soil Science and Agriculture Chemistry, Faculty of Agriculture, SKUAST-J, Chatha, Jammu- 180009, UT of J&K, India.²Division of Fruit Science, Faculty of Horticulture and Forestry, SKUAST-J, Chatha, Jammu- 180009, UT of J&K, India.³Division of Agronomy, Faculty of Agriculture, SKUAST-J, Chatha, Jammu- 180009, UT of J&K, India.⁴Division of Plant Physiology, Faculty of Basic Sciences, SKUAST-J, Chatha, Jammu- 180009, UT of J&K, India.**ABSTRACT**

Climate change, water scarcity, population growth, rising food prices, and other socioeconomic repercussions are anticipated to pose a danger to agriculture and global food security in the twenty-first century, especially for the world's poorest residents of arid and subarid countries. We must concentrate on dry lands to significantly boost the production of grains because agricultural fields with irrigation systems have been fully utilized. Utilizing arid areas to produce sufficient quality grains is difficult because of their poor productivity. The most important source of food in the world is cereal grains, which also play an important part in the global diet of people. In the semiarid tropics of Africa and Asia, millet is one of the most significant drought-resistant crops and is a major source of carbohydrates and proteins for the local population. Additionally, millet grain is currently attracting more interest from food scientists, technologists, and nutritionists due to its significant contribution to national food security and potential health advantages. The phytochemicals found in millet grains have a beneficial impact on human health by reducing phytates and cholesterol levels. The quest for substitute grains is necessary to relieve the pressure caused by the frenetic demands on cereals and their uses in numerous industries. The performance improved when pearl and finger millets were substituted for maize in the diets of several animals. In terms of poor growth conditions and high nutritional value, millets outperform other grains like wheat and rice as crops that are climate change compliant. These strategies will aid in the fight against hunger and malnutrition while also providing monetary benefits to the millet growers and other stakeholders.

Keywords: Climate change, cholesterol, drought tolerant, food security, grains, malnutrition, millets, nutritional value.

1. Introduction

The world's population is projected to grow to 9.1 billion people by 2050 [25]. Urgent action is required to ensure increased food supply and food security considering the restricted land resources. Food security has grown more important on both the international and domestic fronts as a result of the global supply, economic growth, and rising population in emerging countries. Two of the largest difficulties facing the modern world are addressing hunger and feeding the planet's population. Conflicts, insufficient production of food resulting in imbalances in supply and demand, and shortages in the supply of micro- and macronutrients are some of the causes of this problem causing global instability in several places. The concept of food security encompasses both sustained access to wholesome nutrition as well as consistent access to physical food availability. Food insecurity is first divided into two categories by the Food and Agriculture Organisation (FAO): moderate (limited access to food results in poorer dietary quality, eating patterns, health, and wellbeing) and severe (running out of food, experiencing hunger, and having health and wellbeing at severe risk) security [28]. The threat of climate change and global warming persists even while some of these causes of hunger can be addressed,

resulting in a marginal decrease in the number of people suffering from hunger and malnutrition from about one billion in 1990-1992 to 850 million in 2010-2012 [29]. Estimates indicated that by 2050, 2-3 billion people may experience hunger and other food and nutritional insecurities due to the decline in food production rates and the additional burden of feeding a population greater than 9 billion [91]. Crop yields, crop production, and the overall sustainability of our food systems are all said to be directly impacted by climate change and rising world average temperatures. At both the macro and micro levels, ensuring food and nutrition security for India's enormous and diverse population is a challenge, and work has been done on numerous fronts. There is no question that significant efforts have been undertaken recently to achieve food and nutrition security in India, and the advancements made cannot be disputed. However, unless concerted efforts are made, food insecurity and malnutrition appear to be continuing and may continue shortly. So, the goal of sustainable improvement in food and nutrition security continues to be the main objective. According to some estimates, a few places may gain from climate change because of higher production and yields, but this won't be enough to feed the growing population around the world [21]. Furthermore, the majority of scientists concur that crop output would be severely decreased by the current rates of global warming and greenhouse gas emissions. Therefore, ensuring food security depends greatly on reducing greenhouse gas emissions to control global temperatures. However, one of the main sources of greenhouse gases like methane in the environment is the agriculture industry. Intensive agricultural practices, which are practiced in various

*Corresponding Author: **Nikhil Thakur**Email Address: nikhilthakur1024@gmail.comDOI: <https://doi.org/10.58321/AATCCReview.2023.11.04.421>© 2023 by the authors. The license of AATCC Review. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

parts of the world, typically result in higher emissions [60]. In addition to being a large source of macronutrients including carbohydrates, lipids, and proteins, cereal crops also have a sizable potential to contribute to global warming. Wheat has the biggest global warming potential of all the major cereal crops, with an estimated 4 tonnes CO₂ eq/ha, followed by rice and maize (an estimated 3.4 tonnes CO₂ eq/ha). Additionally, the carbon equivalent emissions from these crops are considerable, with an average of 1000, 956, and 935 kg C/ha for wheat, rice, and maize, respectively [43]. They are widely cultivated and the main sources of nutrition for the entire world's population despite having greater emission rates. Other minor cereal crops, including millet and sorghum, have significantly lower carbon footprints. This is one of the primary reasons why millets could be one of the crops that reduce the global carbon footprint [65]. Improved nutritional security can also be achieved by diversifying the agricultural system and using drought-resistant crops. A class of coarse grain cereals known as millets can be very helpful in achieving this. Millets are a group of diverse small-grain cereal crops grown belonging to the Poaceae grass family and are considered one of the oldest cultivated crops. They are composed of up to a dozen crop species, mostly from developing nations in Asia and Africa. Despite significant pressure from competing crops in terms of regulations and production support, India is the world's largest producer of millets, generating approximately 40% of the millets consumed worldwide. Ninety-seven percent of millet is produced in developing nations, making millet significant crops in the semiarid tropical regions of Asia and Africa [88]. In temperate, subtropical, and tropical latitudes, the crop is primarily farmed on marginal lands in arid locations. The crop is preferred because of its yield and brief growth season in hot, dry conditions. Throughout human history, millet has played a significant role as a staple meal, especially in Asia and Africa. Since millets are C₄ plants, they have a much more efficient photosynthetic system than C₃ plants like rice and wheat. The majority of millet grains are now commonly referred to as "Nutri-cereals" because they have higher protein, fiber, calcium, and mineral contents than the widely consumed grains of rice

and wheat. For the past 10,000 years, East Asia has been cultivating them. Millets are native to many regions of the world, and pearl millet is the variety that is grown the most widely. It is a significant crop in India and several regions of Africa [52]. The most significant varieties include foxtail, finger, proso, and pearl millet. Pearl millet accounts for roughly half of the world's millet production. The most significant species of millet in terms of cultivated land and contributions to food security in areas of Africa and Asia is pearl millet. In developed nations and some regions of Asia, proso millet is utilized for food and as bird seed. The crop of foxtail millet is significant in China and Eastern Europe [80]. In the colder, higher-altitude regions of Africa and Asia, finger millet is widely grown as a food crop and used as a popular ingredient in traditional beer. The other species—barnyard, kodo, and little millets, as well as fonios and teff—are regionally or internationally significant food grains. The physical traits, qualitative characteristics, soil, climate requirements, and growth cycles of the various species vary. The scientific and common names of various millets vary depending on the place in which they are grown, and these millets are grown in various parts of the world and require different growth conditions (Table 1). Millions of people every day consume rice, wheat, maize, and, to a lesser extent, millet as their primary sources of nourishment. The growth pattern of these crops is determined by temperature and water availability [66]. While wheat is mostly grown in regions that have restricted water supplies and suitable temperatures, rice, and maize are grown where there is a plentiful supply of water. Sorghum and millets are cultivated in regions with limited water supplies. Due to its resilience to biotic and abiotic challenges and its large production on low-quality lands with little input, millets can even be grown in semi-arid and arid areas [5]. Millets could be crucial functional foods for the prevention of non-communicable diseases because they are a rich source of antioxidants, fiber, and essential and non-essential amino acids, vitamins, and minerals. This review aims to provide detailed provides detailed nutritional composition and its benefits to humans and livestock, the need for growing millets, and various constraints and perspectives associated with it.

Table 1. Scientific names, common names, and growing conditions are required for the cultivation of different types of Millet around the world.

Type of Millets	Scientific Name	Common Name	Growing Conditions	References
Pearl Millet	<i>Pennisetum typhoides</i>	Bulrush millet	Dry climates, marginal soils, Rainfall 200–500 mm	Guigaz and Del'agronome, 2002
Finger Millet	<i>Eleusine corona</i>	Birds food millet or African millet	Resist higher temperatures and salt, with a soil pH range of 5 to 8.2 and a moderate amount of rainfall.	Devi et al., 2014
Porso Millet	<i>Panicum miliaceum</i>	Common millet, hog millet, broom corn, yellow hog, Hershey and white millet	Less water, Rainfall less than 600 mm, average temperature 17 °C during daytime	Zarnkow et al., 2009
Foxtail Millet	<i>Setaria italic</i>	Italian millet, German millet, or hay millet	Less water, short duration	Vetriventhan et al., 2012
Barnyard Millet	<i>Echinochloa crusgalli var. Frumentacea</i>		Drought tolerant, rapid maturation rate	Zhang et al., 2007
Kodo Millet	<i>Paspalum scorbiculatum</i>		High drought resistance, Good yields, period of 80–135 days	Arendt and Dal Bello, 2011

2. Nutritional Importance

There are plenty of chronic diseases and health problems in the world today. According to the 2016 Global Nutrition Report, in 129 nations (countries with data available), 44% of the population suffers from very serious levels of undernutrition, adult overweight, and obesity [42]. The majority of these disorders are caused by nutrient imbalances in the diet. United Nations Food and Agriculture Organisation estimates that 7.9% of the world's population, or 795 million people, were undernourished in 2015. However, more than 1.9 billion adults under the age of 18 (or 39% of the world's population) were reported to be overweight, and another 13% to be obese [26]. The World Health Organisation has previously classified obesity-related problems like diabetes and cardiovascular disease as an epidemic [92]. The majority of undernourished people in the world reside in India. Globally, 4 69,000 deaths were attributed to protein energy malnutrition (PEM), while 84,000 deaths were attributed to a lack of other essential nutrients such as iron, iodine, and vitamin A [89]. With a prevalence incidence of 11% for men and 15% for women, obesity is a significant public health issue in India. Because millets are a great source of several essential elements, they offer an extra benefit in the fight against nutrient deficiencies in third-world countries.

Millets are equivalent to and occasionally have higher nutrient levels than conventional cereals in terms of calorie value, protein, and macronutrients. As a result of their high quantities of calories, calcium, iron, zinc, lipids, and high-quality proteins, they significantly contribute to the diets of both humans and animals. They are also abundant providers of vitamins and dietary fiber. Here, is the nutrient composition of millets:

I. Carbohydrates: The carbohydrates in pearl millet grains include dietary fiber, starch, and soluble sugars. The endosperm of pearl millet contains glucose in the form of amylose and amylopectin and is regarded as having a high starch content. The starch content of different pearl millet grain genotypes varies from 62.8 to 70.5% and approximately 71.82 to 81.02%. Soluble sugars range from 1.2 to 2.6%, and amylose from 21.9 to 28.8% and amylopectin from 21.9 to 28.8%. The starch included in pearl millet can be employed as bulking, thickening, and gelling agents for food texture [35]. However, according to [7], the total carbohydrate content of finger millet ranges from 72 to 79.5%. Additionally, [90] noted that the detailed profile of the carbohydrates ranged between 59.5 and 61.2% for starch, 6.2-7.2% for pentosans, 1.4-1.8% for cellulose, and 0.04-0.6% for lignin.

II. Protein: Protein is the second main component of millet. Pearl millet has 11.6% protein, which is more than the 7.2% in rice, 11.5% in barley, 11.1% in maize, and 10.4% in sorghum [44]. Additionally, [3] noted that pearl millet contained 9.79% protein. The crude protein content of pearl millet grain is thought to be 8–60% higher by weight compared to maize [8]. Contrarily, finger millet has about 5-8% protein [12] recorded the greatest protein level for finger millet at nearly 11%, and [3] documented a proportion of 6.32% in finger millet. Furthermore, [83] revealed that the amino acid profile of pearl millet is equivalent to that of wheat, barley, and rice but contains more lysine, threonine, methionine, and cysteine than sorghum and maize proteins. Moreover, the distribution of proteins in pearl millet grain is thought to be similar to that of maize, notably true prolamins, which are thought to be soluble in alcohol. Additionally, [55] noted a high degree of critical amino

acid balance and endorsed pearl millet as a crucial source of protein and energy for people. Additionally, arginine, threonine, valine, isoleucine, and leucine were discovered to have greater digestibilities in pearl millet than maize. Because finger millet contains more lysine, threonine, and valine than other millet, it is somewhat balanced in terms of its concentration of important amino acids [69].

III. Dietary fiber: Fibre is thought to be beneficial for gut health, and a moderate diet of foods high in fiber may help to promote gut health [55]. Fibre plays a similar role in the prevention of diabetes, colon cancer, and heart disease [24]. Bowel movement is enhanced by the pearl millet's high dietary fiber content, which ranges from 8% to 9% [70]. Additionally, due to its slow digestion, it prolongs the transit time, reducing the absorption of glucose into the blood, which benefits non-insulin-dependent diabetic patients. People who eat millet have a lower incidence of diabetes [73]. Additionally, the fiber in millet may help lower bad cholesterol while raising good cholesterol. It inhibits the body from producing bile acids, which result in gallstones [74]. In addition, pearl millet's high fiber content slows the passage of food from the stomach to the intestines. This results in longer gaps between meals, which helps avoid obesity.

IV. Lipids: The fat content of pearl millet is thought to be between 5 and 7 percent [33] as compared to 3.21 to 7.71 percent in maize [84]. The percentage of lipids in finger and pearl millet was also reported by [40] to be 1% and 5%, respectively. Pearl millet has a high concentration of fatty acids such as palmitic, stearic, and linoleic acids, while maize has a lower percentage of oleic acid [1]. The overall lipid content of pearl millet grains is larger than that of other millet species, ranging from 1.5 to 6.8% [83]. The contents of free and bound lipids in pearl millet are, respectively, 5.6 to 6.1 and 0.6 to 0.9%. [82] reported that the overall lipid content of finger millet was 5.2% (2.2% free lipids, 2.4% bound lipids, and 0.6% structural lipids). On the other hand, oleic, palmitic, and linoleic acids were found to be the main fatty acids found in finger millet. Unsaturated fatty acids make for 74.4% of finger millet's total fatty acid production compared to saturated fatty acids 25.6% [46].

V. Macronutrients: [53], environmental stresses such as high salt concentrations, limited water accessibility, and extreme temperatures have been shown to affect the mineral content of food. The overall mineral and trace element content of pearl millet depends on the type of soil. Ash concentration in pearl millet and maize, respectively, ranges from 1.6 to 3.6% and 0.86 to 1.35%. Pearl millet has higher quantities of minerals than maize, including calcium, phosphorus, magnesium, manganese, zinc, iron, and copper [1]. Due to its high oil content, pearl millet is also regarded as an excellent source of fat-soluble vitamin E (2 mg/100 g). The grain is regarded as an excellent source of vitamin A [83]. [27], pearl millet contains between 45.6 and 48.6 mg of calcium per 100 g. Additionally, it contains a lot of phosphorus, a crucial component found in the mineral matrix of bone, as well as adenosine triphosphate, or ATP, the body's primary energy source. Also, it promotes bone development and healing. According to certain research [58] the bioavailability of iron may be hampered by phytates, oxalates, and polyphenols, which are found in high concentrations in pearl millet. Contrarily, finger millet has a high calcium content that varies from 162 mg/100 g to 487 mg/100 g depending on the genotype

[86]. According to a report by [78], finger millet has 344 mg of calcium per 100 g. Additionally, millet contains a significant amount of magnesium, which is thought to help the body fight diseases like cancer. [6] stated that magnesium content in finger millet ranged from 84.71 mg/100 g to 567.45 mg/100 g. Additionally, finger millet is regarded as a rich source of natural calcium, which promotes bone growth and lowers the incidence of fractures.

Table 2. Nutrient composition of millets in comparison to fine cereals (per 100 g) [36]

Food grain	Carbohydrates (g)	Protein (g)	Fat (g)	Energy (KCal)	Crude fiber (g)	Mineral matter (g)	Calcium (mg)	Phosphorus (mg)	Iron (mg)
Sorghum	72.6	10.4	1.9	349	1.6	1.6	25	222	4.1
Pearl millet	67.5	11.6	5.0	361	1.2	2.3	42	296	8.0
Finger millet	72.0	7.3	1.3	328	3.6	2.7	344	283	3.9
Little millet	67.0	7.7	4.7	341	7.6	1.5	17	220	9.3
Kodo millet	65.9	8.3	1.4	309	9.0	2.6	27	188	0.5
Foxtail millet	60.9	12.3	4.3	331	8.0	3.3	31	290	2.8
Barnyard millet	65.5	6.2	2.2	307	9.8	4.4	20	280	5.0
Proso millet	70.4	12.5	1.1	341	2.2	1.9	14	206	0.8
Wheat (whole)	71.2	11.8	1.5	346	1.2	1.5	41	306	5.3
Rice (raw, milled)	78.2	6.8	0.5	345	0.2	0.6	10	160	0.7

VI. Polyphenols: The body's immune system is believed to be strengthened by the presence of the primary polyphenols, which are abundant in millet and include phenolic acids and tannins [9]. The enzymatic hydrolysis of complex carbs can be somewhat inhibited by millet phenolics, which also inhibit malt amylase, -glucosidase, and pancreatic amylase, all of which lower postprandial hyperglycemia. Similar to this, ferulic and p-coumaric acids, which are present in whole pearl millet, are thought to have the ability to reduce cancer cells [10]. According to [16], the phenols found in millets have been shown to offer several health benefits, including the ability to serve as an antioxidant, anti-inflammatory, and antiviral. In addition, celiac disease is a genetically predisposed condition brought on by consuming gluten. Millets help to prevent celiac disease by minimizing the discomfort brought on by typical cereal grains that contain gluten because they are gluten-free [71]. Millets contain phenolic acids, which occur (60%) in bound and free forms. The most prevalent phenolics in millets are hydroxycinnamic acids, which are primarily located in the phenolic acids' insoluble-bound regions [49]. Ferulic acid, an antioxidant, is the most prevalent form of hydroxycinnamic acid. Antioxidants are well-known substances that reduce the body's exposure to free radical damage and have anti-inflammatory properties [51]. Additionally, millet grains have been discovered to contain ferulate dimers, which exhibit considerable antioxidant activity [23].

Further, the flavonoids found in millet grains include anthocyanidins, chalcones, amino phenolics, flavanols, flavones, and flavanones. Furthermore, according to [22], proanthocyanidins, also known as condensed tannins, are thought to be present in certain millet cultivars. Significant amounts of tannin are primarily found in colored millet types [54]. This finding was linked to the presence of condensed tannins because they significantly contribute to the color of the grain. However, a high concentration of condensed tannins may negatively impact the bioavailability of minerals and proteins [11].

2.1 Other health benefits: [81] reported that the aqueous extract of foxtail millet (*Setaria italica*) has been shown to have anti-hyperglycaemic and anti-lipidemic effects in streptozotocin-induced diabetic rats. According to the study,

diabetic rats received a dose of 300 mg of *Setaria italica* seed aqueous extract per kilogram (kg) of body weight, which led to a significant decline (70%) in blood glucose levels after 6 hours. As evidence of the hypolipidemic effect of the aqueous extract, they also discovered higher levels of HDL (high-density lipoproteins) cholesterol and lower levels of triglycerides, total LDL (low-density lipoproteins), and VLDL (very low-density lipoproteins) cholesterol in diabetic treated rats compared to diabetic untreated rats. [13] stated that Korean foxtail millet's dietary protein has a positive impact on cholesterol and insulin sensitivity. This experiment showed that rats fed foxtail millet had a substantial decrease in insulin levels. When [47] looked into the impact of consuming millet on lipid levels and C-reactive protein concentration, they discovered that contrary to their earlier studies [11], hyperlipidemic rats fed foxtail millet had lower levels of triglycerides. In rats fed foxtail millet, the levels of C reactive protein, a sign of inflammation, were also shown to be lower. Kodo millet aqueous and ethanolic extracts have been shown to cause a dose-dependent decrease in fasting blood glucose [39].

2.2 Dual use of millets as food and feed:

Due to their abundance of essential minerals like calcium, dietary fiber, polyphenols, and protein, millet grains are regarded as one-of-a-kind crops [16]. Many Asian and African nations rely on them as a primary food supply. A larger portion of the millet produced is mostly used for human use, and a smaller portion is used to make beer, livestock, and bird feed. In some regions of Africa, millet is prepared as a porridge with a thin or thick consistency, whereas in other regions, it is prepared as couscous [59]. The effectiveness of pearl millet as a feed item for poultry production was confirmed by research utilizing the whole grain or crushed grain that was used in chicken diets [14]. The primary goal of pearl millet production in Africa and Asia is thought to be grain, while forage is a significant secondary product used for construction, fire, and animal feed [2]. While all-season millets ensure food security, fodder, nutrition, health, and sustainable livelihood, in contrast to season-specific crops like wheat and rice [72]. Because they are free of gluten and have a higher level of nutritional fiber than rice, pearl millet grains have a great deal of promise as a source of nourishment for people. Additionally, they have the same amount of fat as maize

cereal and more critical amino acids like leucine, isoleucine, and lysine than traditional cereals like wheat and rye [41]. In India, where millet is frequently used, it is converted into the dense bread known as dosa, which is formed from a combination of millet and other grains. Additionally, it is used to make roti, sushi, no-yeast pizza, biscuits, and couscous [19]. Another item manufactured from millet is Madua, a popular finger millet-based beverage in India. Additionally, millet is used to make Oshikundu, a traditional alcoholic or non-alcoholic drink in Namibia [45].

2.3 Monetary benefits of using millet.

The cost of millet is believed to be 40% cheaper than that of maize, making it a cheap and gluten-free cereal [32]. [76] estimated that the trade value of pearl millet was less than or equal to 77.78% of the price of maize grain. Pearl millet grain has more protein per grain than maize, it might allow diets to be formulated without protein supplements, lowering the price of food and feed in the process. Aside from that, millet is less expensive to produce than other cereals like sorghum and maize. For instance, the water use of Brazilian pearl millet cultivars is more efficient than that of sorghum and maize grown in semi-arid regions of the country (56 ± 2.8 kg DM/ha/mm water vs. 45 ± 1.9 kg DM/ha/mm water for sorghum; [75] and 21 ± 2.4 kg DM/ha/mm water for the Brazilian maize cultivars (Dos Santos *et al.*, 2010). For instance, the water use of Brazilian pearl millet cultivars is more efficient than that of sorghum and maize grown in semi-arid regions of the country (56 ± 2.8 kg DM/ha/mm water vs. 45 ± 1.9 kg DM/ha/mm water for sorghum [75] and 21 ± 2.4 kg DM/ha/mm water for the Brazilian maize cultivars; [20]. In an investigation by [32], it was discovered that completely substituting pearl millet for maize in the diet of feedlot cattle was the most cost-effective option. The price of lean and fat cattle, initial weight, final weight, cost of concentrate, cost of roughage, consumption of concentrate, and consumption of roughage were listed as the factors that influenced the financial indicator [77]. Furthermore, it is reasonable to believe that millet's positioning as a grain that competes with maize will shift the supply's balance and release pressure on maize consumption, thus, lowering its prices. A different study by [68] found that the cost of feed needed to create one kilogram of live weight gain in chickens fed on maize was higher than the cost for the same amount of gain in chickens fed on pearl millet, finger millet, and sorghum. According to [56], millet grain feed has the lowest cost per kg of feed and the lowest cost of feed per unit weight gain, making broiler chicken production more affordable and cost-effective [93] found that the annual net profit from using pearl millet as the only feedstock was \$25,175,000 in comparison to \$23,758,000 for maize feedstocks, a difference of almost \$1.4 million.

3. Constraints with food security in developing nations

A condition where people have physical and financial access to wholesome food that satisfies their dietary needs is referred to as food security [94]. The scope of food security problems is frequently reduced to the availability of agricultural products like cattle [37]. The difficulties are thought to be more intricate than simply increasing the supplies. Several elements, including urbanization and accessibility, are among the restrictions listed by [37]. The structure and procedures that govern economies and societies, as well as institutional failings, are frequently mentioned as contributing reasons to food insecurity [63]. [30]

reported that the casual intervention to prevent food insecurity is not always simple, therefore proxies of actions and a deeper comprehension of the various potential pathways are crucial for successful interventions. To overcome these intricate obstacles, multisectoral strategies and planning are needed. [63] stated that the best strategy to address the problems of food and nutrition security is to combine the indicators of food insecurity with the relevant socioeconomic and environmental variables of a specific entity. Despite the difficulties facing food security in developing nations, the importance of locally grown and underutilized grains like millet cannot be understated. Some of these problems cannot be solved by the standard grain crops [57]. The expansion of small grains like millet could worsen poverty among the underprivileged population, especially in light of recent events like the extraordinary COVID-19 epidemic [57]. Millets are crops that, in the context of climatic change, have the potential to endure challenging circumstances and contribute to the stability of food security. [62] reported that minor millets like finger, kodo, foxtail, tiny, porso, and barnyard can flourish in a variety of soils, different climatic conditions, distinct photoperiods, and due to their genetic adaptability. These qualities make millets suitable for replacing staples like wheat and rice in difficult climate zones, eventually resulting in food security in these regions. However, millet is regarded as a neglected agro-biodiversity, even though it can improve the food security of underprivileged people in developing countries [31].

4. Impact of processing on millet grains' antioxidant activity

The quantity and activity of the antioxidants in millet grains were also discovered to be impacted by processing techniques such as malting, decortication, soaking, and boiling. In one study, finger millet was malted for 96 hours, and the antioxidant capacity of the fraction carrying free phenolic acids rose (2-fold), while the antioxidant capacity of the fraction holding bound phenolic acids was decreased [68]. In another study, a growth-promoting medium was created to increase the production of a water-soluble protein from germinated millet that is inhibitory of hydroxyl radicals. [48] reported that the single-factor test showed that H_2O_2 is essential for the inhibitory activity. Additionally, the effects of the sprouting conditions (temperature, duration, and pH of stress media) on the hydroxyl radical inhibition were examined to further boost the yield of the water-soluble protein that inhibits hydroxyl radicals from stress-germinated millet. The ideal parameters were found to be pH 7.5 in the stress medium, a culture duration of 54 hours, and a temperature of $28^\circ C$. The maximum inhibition (60.38%) was attained under ideal circumstances [48]. Additionally, 3 days of foxtail millet germination allowed researchers to produce flour with a strong DPPH-scavenging activity [15].

The antioxidant activity of Kodo or finger millet decreased after roasting or boiling. Kodo millet's DPPH quenching activity was likewise reduced by fractionating it into husk and endosperm, and the phytochemicals appeared to work synergistically with one another [38]. Additionally, the impact of germination, steaming, and roasting on the antioxidant qualities of little millet was also examined. The outcomes revealed that the total phenolic, flavonoid, and tannin levels of little millet increased by 21.2, 25.5, and 18.9 mg/100 g, respectively, over the original sample [64]. Dehulling and hydrothermal treatments, however, were discovered to have an impact on the phenolic content and antioxidant potential of pearl millet grains [9]. Due to oxidation and degradation of chemicals that occur during heat treatments

like cooking, boiling, and roasting, antioxidant levels and activity have decreased. However, the removal of the pericarp layer from the grains, which is known to be high in polyphenol and antioxidant chemicals, can be blamed for the decline that resulted from dehulling. As a result, millet grains, their fractions, and food products must be processed under ideal circumstances to preserve their quality and possible health advantages. Additionally, the endogenous enzymes that are converted to simpler molecules with enhanced antioxidant activity during germination are responsible for the rise in antioxidant levels and their activities.

5. Challenges and future perspectives

The data reviewed above shows that despite the nutritious richness and potential health advantages of millet grains, were comparable to popular grains like wheat, rice, and maize, and although processing techniques like fermentation, soaking, malting, and fortification/supplementation were discovered to improve its edibility and nutritional qualities, millet grain consumption is still primarily restricted to rural communities at the household level. This is a result of the absence of cutting-edge millet processing technologies that can produce safe, easy-to-handle, ready-to-cook or ready-to-eat goods and meals on a big scale that can be utilized to feed numerous people in metropolitan areas [85]. Nevertheless, as the world's population grows, so do the demands for food, fuel, and other necessities. As a result, society will be under pressure to either change its current crop consumption habits or increase agricultural production [50]. Along with increased yields, crop diversification must also be promoted at the national and household levels. One crucial element of therapeutic dietary modification and encouraging consumption of minor-grain foods is the provision of more traditional and nutritious whole-grain and multigrain alternatives for refined carbohydrates [78]. The role of gluten protein in generating high-quality, manageable baked goods and other grain-based dishes that demand elastic and extensible dough is well established. However, considering that millet grains are gluten-free and based on the outcomes of several laboratory tests, it appears that they are not ideal for making pure-millet bread and other solid food products that are simple to handle. To encourage the use of millet grains, it appears that using it as a substitute for wheat in composite flours, complementary foods, and food blends is the best way to create nutrient-dense, "healthy," and secure food products that are of high caliber and can be stored for a long time. Additionally, there is a need for cutting-edge processing technology for decortication, milling, and other preparation procedures of millet grain food to make high-quality products on a commercial scale for urban consumers. In return, it is necessary to produce millet cultivars with high essential amino acid content and maintain a steady supply of premium millet grains for industrial needs. Future research studies should assess the nutritional value and potential health advantages of millet grains and their fractions in animal and human models to support efforts to promote their use as food.

6. Conclusion

Millet grains were shown to offer a significant potential to improve food and nutritional security. Millets should therefore be added to the list of staple foods together with wheat and rice. Millets may grow well in adverse conditions like drought, and some wild varieties can even thrive in wetlands and populated areas. These contain gluten-free protein, low glycemic index

carbohydrates, and an abundance of minerals (such as calcium, iron, copper, magnesium, and others), B vitamins, and antioxidants. These exceptional qualities make them nutrient-dense and resistant to climate change crops. These can benefit the health of the community as a whole in addition to providing farmers with an additional source of revenue. As a result, research and development initiatives as well as the creation of policies are needed; some actions have already been done globally, particularly in India and some are required to be taken. Minor millets have received very little attention in the scientific community despite their great nutritional qualities and easy production methods. They are frequently disregarded, leading to the term "orphan cereals" being used by scientists to describe them. However, to increase the micronutrients' bioavailability and raise the caliber of millet diets, new processing and preparation techniques are required. The bioavailability, metabolism, and health benefits of millet grains and their many components in humans require further study. The current nutrient deficiencies of protein, calcium, and iron in poor nations will be addressed by the inclusion of millet-based foods in international, national, and state-level feeding programs.

Conflict of interest: The authors declare no conflict of interest.

Acknowledgment: The support provided by all the authors has been duly acknowledged.

7. References

1. Adeola O, Orban JI. Chemical composition and nutrient digestibility of pearl millet (*Pennisetum glaucum*) fed to growing pigs. *J Cereal Sci.* 1995;22:177–84.
2. Andrews DJ, Kumar KA. Pearl millet for food, feed, and forage. *Adv Agron.* 1992;48:89–139.
3. Anitha S, Govindaraj M, Kane-Potaka J. Balanced amino acid and higher micronutrients in millets complements legumes for improved human dietary nutrition. *Cereal Chem.* 2019;97:74–84.
4. Arendt, E.; Dal Bello, F. *Gluten-Free Cereal Products and Beverages*; Academic Press: Cambridge, MA, USA, 2011.
5. Awika, J.M. Major cereal grains production and use around the world. In *Advances in Cereal Science: Implications to Food Processing and Health Promotion*; ACS Publications: Washington, DC, USA, 2011; pp. 1–13.
6. Bachar K, Mansour E, Ben Khaled A, Abid M, Haddad M, Ben Yahya L, Jarray EL, N, Ferchichi A. Fiber content and mineral composition of the finger millet of the oasis of Gabes Tunisia. *J Agric Sci.* 2013.
7. Bhatt A, Singh V, Shrotria PK, Baskheti DC. Coarse grains of Uttaranchal: ensuring sustainable food and nutritional security. *Indian Farmer's Digest.* 2003;7:34–8.
8. Burton GW, Wallance AT, Radice KO. Chemical composition during maturation and nutritive value of pearl millet. *Crop Sci.* 1972;12:187–92.

9. Chandrasekara A, Shahidi F. 2012. Bioaccessibility and antioxidant potential of millet grain phenolics as affected by simulated in vitro digestion and microbial fermentation. *J Funct Foods* 4:226–37.
10. Chandrasekara A, Shahidi F. Determination of antioxidant activity in free and hydrolyzed fractions of millet grains and characterization of their phenolic profiles by HPLC-DAD-ESI-MSn. *J Funct Foods*. 2011;3:144–58.
11. Chavan UD, Shahidi F, Naczki M. Extraction of condensed tannins from beach pea (*Lathyrus maritimus* L.) as affected by different solvents. *Food Chem*. 2001;75:509–12.
12. Chethan S, Malleshi NG. Finger millet polyphenols: optimization of extraction and the effect of pH on their stability. *Food Chem*. 2007;105:862–70.
13. Choi YY, Osada Y, Ito, Nagasawa T, Choi MR, Nishizawa N. Effects of dietary protein of Korean foxtail millet on plasma adiponectin, HDL cholesterol, and insulin levels in genetically type 2 diabetic mice. *Biosci Biotechnol Biochem*. 2005;69(1):31–7.
14. Cisse RS, Hamburg JD, Freeman ME, Davis AJ. Using locally produced millet as a feed ingredient for poultry production in Sub-Saharan Africa. *J Appl Poultry Res*. 2016;26(1):9–22. <https://doi.org/10.3382/japr/pfw042>.
15. Coulibaly A, Chen J. 2011. Evaluation of energetic compounds, antioxidant capacity, some vitamins and minerals, phytase and amylase activity during germination of foxtail millet. *Am J Food Technol* 6(1):40–51.
16. Devi PR, Vijayabharathi R, Sathyabama S, Malleshi NG, Priyadarisini VB. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. *J Food Sci Technol*. 2014;6:1021–40.
17. Devi PR, Vijayabharathi R, Sathyabama S, Malleshi NG, Priyadarisini VB. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. *J Food Sci Technol*. 2014;6:1021–40. <https://doi.org/10.1007/s13197-011-0584-9>.
18. Devi, P.B.; Vijayabharathi, R.; Sathyabama, S.; Malleshi, N.G.; Priyadarisini, V.B. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: A review. *J. Food Sci. Technol*. 2014, 51, 1021–1040.
19. Dias-Martins AM, Kênia Letícia P, Sidney P, José Avelino SR, Carlos WPC. Potential use of pearl millet (*Pennisetum glaucum* (L.) R. Br.) in Brazil: food security, processing, health benefits and nutritional products. *Food Res Int*. 2018;109:175–86. <https://doi.org/10.1016/j.foodres.2018.04.023>.
20. Dos Santos RD, Pereira LGR, Neves ALA, Azevedo JAG, Demoraes SA, Costa CTF. Agronomic characteristics of maize varieties for silage production in the submédio São Francisco river valley. *Acta Sci Anim Sci*. 2010;32:367–73.
21. Downing, T.E.; Barrow, E.M.; Brooks, R.J.; Butterfield, R.E.; Carter, T.R.; Harrison, P.A.; Hulme, M.; Olesen, J.E.; Porter, J.R.; Schellberg, J.; et al. Quantification of uncertainty in climate change impact assessment. In *Climate Change, Climatic Variability and Agriculture in Europe*; Environmental Change Institute: Oxford, UK, 2000; pp. 415–434.
22. Dykes L, Rooney LW. Sorghum and millet phenols and antioxidants. *J Cereal Sci*. 2006;44(3):236–51.
23. Emiola LO, De la Rosa LC. Characterization of Pearl Millet Non-starchy Polysaccharides. *J Food Sci*. 1981;46:781–5.
24. Eshak ES, Iso H, Date C, Kikuchi S, Watanabe Y, Wada Y, Wakai K, Tamakoshi A. Dietary fiber intake is associated with reduced risk of mortality from cardiovascular disease among Japanese men and women. *J Nutr*. 2010;140:1445–53.
25. FAO, 2009a. *Global Agriculture Towards 2050*. FAO, Rome.
26. FAO. *The State of Food Insecurity in the World 2015*. Food and Agricultural Organization of the United Nations; 2015. <http://www.fao.org/worldfoodsituation/csdb/en/>.
27. Florence SP, Asna U, Asha MR, Jyotsna R. Sensory, physical, and nutritional qualities of cookies prepared from pearl millet (*Pennisetum typhoideum*). *J Food Processing Technol*. 2014;5(10):377.
28. Food and Agriculture Organization 2019. *The state of food security and nutrition in the world: Safeguarding against economic slowdowns and downturns*. Rome, Italy.
29. Food and Agriculture Organization of the United Nations. *Technical Note: FAO Methodology to Estimate the Prevalence of Undernourishment*; FAO: Rome, Italy, 2012.
30. Fraval S, Yameogo V, Ayantunde A, et al. Food security in rural Burkina Faso: the importance of consumption of own-farm sourced food versus purchased food. *Agric & Food Secur*. 2020;9:2. <https://doi.org/10.1186/s40066-020-0255-z>.
31. Garí, JA. Review of the African millet diversity http://www.fao.org/fleadmin/templates/esw/esw_new/documents/Links/publications_other/6_millet.pdf. Accessed 23 Nov 2020.
32. Gomes PC, Rodrigues MP, Albino LFT, Rostagno HS, Gomes MFM, De Mello HH, Brumano G. Determination of chemical composition and energy value of millet and their use in rations of broilers from 1 to 21 days of age. *Rev Bras Zootec*. 2008;37(9):1617–21. <https://doi.org/10.1590/S1516-35982008000900013>.
33. Gopalan C, Rama Sastri BV, Balasubramanian SC. *Nutritive value of Indian foods*. Hyderabad: National Institute of Nutrition; 2003.
34. Guigaz, M. *Memento Del'agronome*; CIRAD-GRET and Ministère des Affaires Étrangères: Montpellier, France, 2002.

35. Hadimani NA, Muralikrishna G, Tharanathan RN, Malleshi NG. Nature of carbohydrates and proteins in three pearl millet cultivars varying in processing characteristics and kernel texture. *J Cereal Sci.* 2001;33:17–25.
36. Hariprasanna, K., Gomashe, S., Ganapathy, K.N. and Patil, J.V., 2014. Millets for ensuring nutritional security.
37. Hatab AA, Cavinato MER, Lagerkvist CJ. Urbanization, livestock systems and food security in developing countries: a systematic review of the literature. *Food Secur.* 2019;11(2):279–99.
38. Hegde PS, Chandra TS. 2005. ESR spectroscopic study reveals higher free radical quenching potential in kodo millet (*Paspalum scrobiculatum*) compared to other millets. *Food Chem* 92:177–82.
39. Hegde PS, Chandrakasan G, Chandra TS. Inhibition of collagen glycation and crosslinking in vitro by methanolic extracts of finger millet (*Eleusine coracana*) and kodo millet (*Paspalum scrobiculatum*). *J Nutr Biochem.* 2002;13:517–21.
40. Himanshu K, Chauhan M, Sonawane SK, Arya SS. Nutritional and nutraceutical properties of millets: a review. *Clin J Nutr Diet.* 2018;1(1):1–10.
41. ICRISAT. Smart food millet recipes 2016. <http://www.icrisat.org/PDF/Food-Booklet-Millet.pdf>. Accessed 16 May 2020.
42. IFPRI. Global Nutrition Report: Malnutrition Becoming the “New Normal” Across the Globe; 2016. <https://www.ifpri.org/news-release/global-nutrition-report-malnutrition-becoming-%E2%80%9Cnewnormal%E2%80%9D-across-globe>.
43. Jain, N.; Arora, P.; Tomer, R.; Mishra, S.V.; Bhatia, A.; Pathak, H.; Chakraborty, D.; Kumar, V.; Dubey, D.; Harit, R. Greenhouse gases emission from soils under major crops in northwest India. *Sci. Total Environ.* 2016, 542, 551–561.
44. Jha A, Tripathi AD, Alam T, Yadav R. Process optimization for manufacture of pearl millet-based dairy dessert by using response surface methodology (RSM). *J Food Sci Technol.* 2013;50(2):367–73.
45. Kumar A, Tomer V, Kaur A, Kumar V, Gupta K. Millets: a solution to agrarian and nutritional challenges. *Agric Food Secur.* 2018;7:31. <https://doi.org/10.1186/s40066-018-0183-3>.
46. Kunyanga CN, Imungi JK, Velingiri V. Nutritional evaluation of indigenous foods with potential food-based solution to alleviate hunger and malnutrition in Kenya. *J Appl Biosci.* 2013;67:5277–88.
47. Lee SH, Chung IM, Cha YS, Park Y. Millet consumption decreased serum concentration of triglyceride and C-reactive protein but not oxidative status in hyperlipidemic rats. *Nutri Res.* 2010;30(4):290–6.
48. Li J, Chen Z, Yao H, Xu Y. 2007. Optimization of stress medium enhance hydroxyl radical inhibition by water-soluble protein from germinated millet. *LWT—Food Sci Technol* 40:1630–6.
49. Liang S, Liang K. Millet grain as a candidate antioxidant food resource: a review. *Int J Food Prop.* 2019;22(1):1652–61.
50. Licker R, Johnston M, Foley JA, Barford C, Kucharik CJ, Monfreda C, Ramankutty N. 2010. Mind the gap: how do climate and agricultural management explain the 'yield gap' of croplands around the world? *Global Ecol Biogeogr* 19:769–82.
51. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev.* 2010;4(8):118–26.
52. Lu H, Zhang J, Liu KB, Wu N, Li Y, Zhou K, et al. Earliest domestication of common millet (*Panicum miliaceum*) in East Asia extended to 10,000 years ago. *Proc Natl Acad Sci USA* 2009;106(18):736772.
53. Martínez-Ballesta MC, Dominguez-Perles R, Moreno DA, Muries B, Alcaraz-López C et al. Minerals in plant food: effect of agricultural practices and role in human health. A review. *Agronomy for Sustainable Development*, Springer Verlag/EDP Sciences/INRA, 2010. Doi: <https://doi.org/10.1051/agro/2009022>.
54. McDonough CM, Rooney LW, Serna-Saldivar SO. In: Kurl K, Ponte JG Jr, editors. The millets in handbook of cereal science and technology, second edition, revised and expanded, chap 4. New York: Marcel Dekker Inc; 2000.
55. McIntosh GM, Noakes M, Royle PJ, Foster PR. Whole-grain rye and wheat foods and markers of bowel health in overweight middle-aged men. *Am J Clin Nutr.* 2003;77:967–74.
56. Medugu CI, Kwari ID, Igwebuike J, Nkama I, Mohammed ID, Hamaker B. Performance and economics of production of broiler chickens fed sorghum or millet as replacement for maize in the semi-arid zone of Nigeria. *Agric Biol J North Am.* 2010;1:321–5.
57. Muthamilarasan M, Prasad M. small millets for enduring food security amidst pandemics. <https://doi.org/https://doi.org/10.1016/j.tplan ts.2020.08.008>. Accessed 23 Nov 2020
58. Nambiar VS, Dhaduk JJ, Sareen N, Shahu T, Desai R. Potential functional implications of pearl millet (*Pennisetum glaucum*) in health and disease. *J Appl Pharm Sci.* 2011;10:62–7.
59. Obiana AB. Overview: importance of millets in Africa 2003. Published online at <http://www.afripro.org.uk/papers/Paper02Obil>. Accessed 7 Feb 2020.
60. Olesen, J.E.; Bindi, M. Consequences of climate change for European agricultural productivity, land use and policy. *Eur. J. Agron.* 2002, 16, 239–262.

61. Ouattara-Cheik AT, Aly S, Yaya B, Alfred TS. A comparative study on nutritional and technological quality of fourteen cultivars of pearl millets *Pennisetum glaucum* (L) Leek in Burkina Faso. *Pakistan J Nutr.* 2006;5(6):512-21.
62. Padulosi S, Mal B, Ravi SB, Gowda J, Gowda KTK, Shanthakumar G, Yenagi N, Dutta M. Food security and climate change: role of plant genetic resources of minor millets. *Indian J Plant Genet Resour.* 2009;22(1):1-16.
63. Pangaribowo, EH, Gerber, N, & Torero, M. Food and nutrition security indicators: a review. *FOODSECURE working paper 04.* 2013
64. Pradeep SR, Guha G. 2011. Effect of processing methods on the nutraceutical and antioxidant properties of little millet (*Panicum sumatrense*) extracts. *Food Chem* 126:1643-7.
65. Prasad, P.V.; Staggenborg, S.A. Growth and production of sorghum and millets. In *Soils, Plant Growth and Crop Production*; EOLSS Publishers Co., Ltd.: Oxford, UK, 2009; Volume 2.
66. Prasad, P.V.; Staggenborg, S.A. Growth and production of sorghum and millets. In *Soils, Plant Growth and Crop Production*; EOLSS Publishers Co., Ltd.: Oxford, UK, 2009; Volume 2. 9.
67. Rama Rao SVG, Shyam Sundar AK, Panda MR, Reddy MV, Raju LN, Praharaj NK. Utilization of different millets replacing maize in coloured broiler chicken diet. *Indian J Anim Nutr.* 2002;19(4):353-8.
68. Rao MVSSTS, Muralikrishna G. 2002. Evaluation of the antioxidant properties of free and bound phenolic acids from native and malted finger millet (ragi, *Eleusine coracana* Indaf-15). *J Agric Food Chem* 50(4):889-92.
69. Ravindran G. Studies on millets: proximate composition, mineral composition, and phytate and oxalate contents. *Food Chem.* 1991;39(1):99-107.
70. Rooney LW, Miller FR. Variation in the structure and kernel characteristics of sorghum. In: *proceeding of the international symposium on sorghum grain quality.* iCRISAT. 28-31. Patancheru, India. 1982; 143-162.
71. Saleh ASM, Zhang Q, Chen J, Shen Q. Millet grains: nutritional quality, processing, and potential health benefits. *Compr Rev Food Sci Food Saf.* 2013;12:281-95.
72. Shivananjappa M. Meats are malady and millets are magical. *J Nutr Food Technol.* 2018;1(1):14-5. <https://doi.org/10.30881/jnfrt.00004>.
73. Shobana S, Sreerama YN, Malleshi NG. Composition and enzyme inhibitory properties of finger millet (*Eleusine coracana* L.) seed coat phenolics: mode of inhibition of α -glucosidase and pancreatic amylase. *Food Chem.* 2009;115(4):1268-73.
74. Shweta M. Pearl millet nutritional value and medicinal uses. *IJARIE-ISSN (O).* 2015;1(3):2395-4396.
75. Silva DA, Santos TC, Azevedo EM, Edvan JAG, Perazzo RL, Pinho AF, Rodrigues RMA, DA SILVA, D. S. Agronomic divergence of sorghum hybrids for silage yield in the semiarid region of Paraíba. *R Bras Zootec.* 2011;2011(40):1886-93.
76. Silva GAH, Restle J, Missio RL, Bilego UO, Fernandes JJR, Rezende PLP, Medeiros Da Silva R, Pereira MLR, Lino FA. Milheto em substituição ao milho na dieta de novilhos confinados. *Semina Ciências Agrárias.* 2014;35:2077-94.
77. Silva RM, João Restle RZ, Fabricio EA, Camera A, Maysonnave GS, Ubirajara O, Bilego UO, Pacheco PS, Vaz FN. Economic analysis of the risk of replacing corn grains (*Zea mays*) with pearl millet grains (*Pennisetum glaucum*) in the diet of feedlot. *Cienc Rural.* 2020. <https://doi.org/10.1590/0103-8478cr20190443>.
78. Singh P, Raghuvanshi RS. 2012. Finger millet for food and nutritional security. *Afr J Food Sci* 6(4):77-84.
79. Singh P, Raghuvanshi RS. Finger millet for food and nutritional security. *African J Food Sci.* 2012;6(4):77-84.
80. Singh, R.B., Khan, S., Chauhan, A.K., Singh, M., Jaglan, P., Yadav, P., Takahashi, T. and Juneja, L.R., 2019. Millets as functional food, a gift from Asia to Western World. In *The Role of Functional Food Security in Global Health* (pp. 457-468). Academic Press.
81. Sireesha Y, Kasetti RB, Nabi SA, Swapna S, Apparao C. Antihyperglycemic and hypolipidemic activities of *Setaria italica* seeds in STZ diabetic rats. *Pathophysiology.* 2011;18:159-64.
82. Sridhar R, Lakshminarayana G. Contents of total lipids and lipid classes and composition of fatty acids in small millets: foxtail (*Setaria italica*), proso (*Panicum miliaceum*), and finger (*Eleusine coracana*). *Cereal Chem.* 1994;71:355-9.
83. Taylor JRN. In: Wrigley C, Corke H, Walker CE, editors. *Millet: in encyclopaedia in grain science*, vol. 2. London: Elsevier; 2004. p. 253-6.
84. Ullah I, Ali M, Farooqi A. Chemical and Nutritional Properties of Some Maize (*Zea mays* L) Varieties Grown in NWFP. *Pakistan P J N.* 2010;9(11):1113-7.
85. Ushakumari SR, Latha S, Malleshi NG. 2004. The functional properties of popped, flaked, extruded and roller-dried foxtail millet (*Setaria italica*). *Intl J Food Sci Technol* 39:907-15.
86. Vadivoo AS, Joseph R, Ganesan NM. Genetic variability and calcium contents in finger millet (*Eleusine coracana* L. Gaertn) in relation to grain colour. *Plant Foods Hum Nutr.* 1998;52(4):353-64.
87. Vetriventhan, M.; Upadhyaya, H.; Anandakumar, C.; Senthilvel, S.; Parzies, H.; Bharathi, A.; Varshney, R.; Gowda, C. Assessing genetic diversity, allelic richness and genetic relationship among races in icrisat foxtail millet core collection. *Plant Genet. Resour.* 2012, 10, 214-223.

88. Vinoth K, Ravidran R. Biofortification in Millets: a sustainable approach for nutritional security front. *Plant Sci* 2017. Available from: <https://doi.org/10.3389/fpls.2017.00029>.
89. Von Grebmer K, Bernstein J, Hossain N, Brown T, Prasai N, Yohannes Y. 2017 global hunger index: The inequalities of hunger. International Food Policy Research Institute. 2017.
90. Wankhede DB, Shehnaj A, Rao MR. Carbohydrate composition of finger millet (*Eleusine coracana*) and foxtail millet (*Setaria italica*). *Plant Foods Hum Nutr.* 1979a;28:293–303.
91. Wheeler, T.; Von Braun, J. Climate change impacts on global food security. *Science* 2013, 341, 508–513.
92. WHO. Obesity and overweight Fact sheet N°311"; 2015. <http://www.who.int/mediacentre/factsheets/fs311/en/>. Accessed 23 Jan 2018.
93. Wilson JP, McAloon AJ, YeeW, McKinneyJ, WangD, BeanSR. Biological and economic feasibility of pearl millet as a feedstock for ethanol production. 2007.
94. World Food Summit. The world free from hunger. Rome; 1996
95. Zarnkow, M.; Mauch, A.; Burberg, F.; Back, W.; Arendt, E.K.; Kreis, S.; Gastl, M. Proso millet (*Panicum miliaceum* L.) a sustainable raw material for the malting and brewing process: A review. *Brew. Sci.* 2009, 62, 119–140.
96. Zhang, J.-P.; Liu, T.-S.; Zhang, J.-P.; Liu, T.-S.; Zheng, J.; Zhang, J.-P.; Liu, T.-S.; Zheng, J.; Jin, Z. Cloning and characterization of a putative 12-oxophytodienoic acid reductase cDNA induced by osmotic stress in roots of foxtail millet: Full length research paper. *DNA Seq.* 2007, 18, 138–144.