

Original Research Article


Open Access

Hygienic collection and preservation of neera from palmyrah palm

 Md. Shamsheer Ahmad¹, Rahul Kumar¹, M.W. Siddiqui¹, Fozia Homa², M. Preema Devi³, and Md. Arshad Anwer^{4*}
¹Food Science and Postharvest Technology, Bihar Agricultural University, Sabour, Bhagalpur, (Bihar)-813210, India

²Department of Statistics, Mathematics and Computer Application, Bihar Agricultural University, Sabour, Bhagalpur, (Bihar)-813210, India

³Department of Pomology and Postharvest Technology, Faculty of Horticulture, UBKV, Pundibari, Cooch Behar (WB)-736165, India

⁴Department of Plant Pathology, Bihar Agricultural University, Sabour, Bhagalpur, (Bihar)-813210, India


ABSTRACT

Palmyrah palm (*Borassus flabellifer* L) produces inflorescence which secretes a sap upon tapping is called neera. In India, palmyrah palm is grown in many states like Andhra Pradesh, Gujarat, Tamil Nadu, Odisha, West Bengal, Bihar, Karnataka, and Maharashtra on large scale. Many challenges are faced by the tappers while tapping neera. The present method of tapping is not scientific. It is highly unhygienic and cross contamination with insects, bacteria and yeast occurs during tapping of neera and fermentation starts quickly after tapping. The fermented product is called toddy which is highly injurious to health. Keeping the problems and challenges of neera tapping and preservation without severe heating, the research investigation entitled "Hygienic collection and preservation of neera from palmyrah palm" has been taken. Application of anti fermenting solution (AFS) of 1500 ppm SO₂ and 0.2% citric acid on the sliced portion followed by collection of neera directly into a polyethylene lined double walled tin container containing 300 g crushed ice and 0.5% lime powder was found effective. The neera was then preserved by thermosonication (35 kHz for 15 and 20 minutes + Pasteurization at 65 °C for 10 minutes). The preserved neera was successfully stored at 2-3 °C for 60 days with stable sensorial quality and without any fermentation.

Keywords: Palmyrah palm, neera, fermentation, hygienic collection, pasteurization, thermosonication and preservation

Introduction

Palmyrah palm (*Borassus flabellifer* L.) belongs to the family Palmae and grown in many countries of the world. The major growing countries are Srilanka, Pakistan, Bangladesh, Myanmar, Thailand, Malaysia, Indonesia, Nigeria, Congo, Sudan, Tanzania, India, China, Jawa, Sulawesi, Cambodia, Laos and Vietnam [1, 2]. In India, palmyrah palm is grown in many states like Andhra Pradesh, Gujarat, Tamil Nadu, Odisha, West Bengal, Bihar, Karnataka, and Maharashtra on large scale. Among Indian states, Tamil Nadu tops the rank as number one producing state with 60% of total production [3]. According to an estimate, India is a home of approximately 102 million palms and 50% of the total population is found in Tamil Nadu [4]. About 12 to 15 years old palmyrah palm produces inflorescence that produces a sweet sap upon tapping is called neera [2]. It is clear in appearance having pleasant aroma, sweet flavor and possess immense medicinal properties beneficial for human health like anti-phlegmatic agent [5], anti-inflammatory, antibacterial, analgesic and have anti-oxidant properties [6]. It Prevents certain types of cancers in human body [7]. The sugar present in neera has low glycemic index (GI-35) which make it safe to drink for diabetic persons [8]. Sugar prepared from neera is reported to have beneficial for curing liver disorder [9]. Many challenges are faced by the tappers while tapping neera. The present method of tapping is not scientific.

It is highly unhygienic. Many cross contamination with insects, bacteria and yeast occurs during tapping of neera and fermentation starts quickly into alcohol (plate 1). A hygienic collection of neera is possible in a polyethylene lined low cost tin box (plate-2) yields fresh neera (plate-3). A front and side view of the same box is shown (plate-4 and 5). Descriptions of plates are mentioned in annexure I.



Plate 1 Unhygienic neera Plate 2 Hygienic collection of neera Plate 3 Fresh neera



Plate 4 Neera collection box (front view) Plate 5 Neera collection box (side view)

*Corresponding Author: **Md. Arshad Anwer**

DOI: <https://doi.org/10.21276/AATCCReview.2025.13.01.302>

© 2025 by the authors. The license of AATCC Review. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Preservation of neera is the biggest challenge faced by palmyrah growers and tappers. It is because of rapid fermentation of neera at ambient temperature. The fermented neera is called toddy. The degree of fermentation of the neera depends on the practice of neera collection and the environmental condition prevailing in the region [10]. Since neera is rich source of sugars, vitamins, proteins and minerals [5], fermentation starts within one or two hours of extraction. A large number of yeast and bacterial strains was found responsible for fermentation of neera [11]. The most common microbial flora found in fresh neera are *Bacillus*, *Lactobacilli*, *Micrococci*, *Enterobacter*, *Leuconostoc*, *Saccharomyces*, *Candida*, and *Pichia* [12]. According to Kurian and Peter, [13], Toddy is the term used to describe the process by which wild yeasts and bacteria ferment sugar sap of neera. The fermented neera (toddy) is a spoiled form of neera. The toddy is, in fact, harmful because it is alcoholic in nature with 5 per cent alcohol and above hence not recommended as a health drink [14]. The business of toddy is prohibited by law in many states of India. It is a total loss for the producers and neera entrepreneurs [15]. Hence collection and preservation of neera by arresting fermentation is challenging and a limiting factor for commercialization of fresh neera without fermentation. Further, any processing or value addition in palmyrah palm is possible only after preservation of neera. Pasteurization is mainly applied for preservation of neera. Mostly pasteurization temperature is kept at boiling temperature. At this temperature, thermal degradation in colour, flavour and nutritional loss is obvious.

Table 1. Treatment Details of experiment I (Hygienic collection of neera)

Treats.	Treatment details
T ₁	No Sanitization, Earthen pot (Labni), Traditional Tapping (Control)
T ₂	Sanitization with AFS* Solution (1500 ppm So ₂ +0.2% C.A.) +Polyethylene (PE) lined labni
T ₃	Sanitization with AFS* Solution (1500 ppm So ₂ +0.2% C.A.) +Polyethylene (PE) lined plastic pot
T ₄	Sanitization with AFS* Solution (1500 ppm So ₂ +0.2% C.A.) +PE lined plastic Pot containing lime (0.5%) + protected by cushioning with thermocol.
T ₅	Sanitization with AFS* Solution (1500 ppm So ₂ +0.2% C.A.) + PE lined plastic Pot containing (1.0%) + protected by cushioning with thermocol+ 200 g Crushed ice
T ₆	Sanitization with AFS* Solution (1500 ppm So ₂ +0.2% C.A.) + Plastic Pot lime (0.5%) + protected by cushioning with thermocol+ 300 g Crushed ice

*AFS- Anti-Fermenting Solution

EXPERIMENT I: Hygienic collection of neera

Under this experiment, neera was collected as per the treatments (table 1). First of all, inflorescence was sliced (plate 6) and sanitized as per treatments by applying an anti fermenting solution on the cut portion as well as on other parts of the inflorescence (plate 7). Now collection container was hanged below the sliced and sanitized inflorescence for collection of neera. The collected neera was brought to the laboratory of the department of Food Science and Postharvest Technology, Bihar Agricultural College, Sabour, Bhagalpur and strained through centrifugation to remove suspended and foreign material (plate 8) and stored at low temperature (0 to 2°C) in order to stabilize the sap and bring it at uniform temperature before initial biochemical analysis.





Plate 6. Sliced Inflorescence for Plate 7. Application of anti Plate 8. Filtered neera tapping neera fermenting solution

The biochemical analysis was performed as per standard procedure. For judging the best tapping method, four parameters were used as mentioned in table 2.

The most common changes upon heating sap is Millard reaction and Caramelisation [16]. Non-thermal processing like ultrasound coupled with low temperature pasteurization called thermosonication can be applied. It has higher efficiency for killing of microbes and deactivation of enzymes at relatively low temperature [17]. Kapturowska et al. [18] also reported that thermosonication is very effective against yeast. Preserved neera can be used in preparation of various value-added products, which may be a source of employment in the palmyra palm growing areas. Keeping the problems and challenges of neera tapping and preservation without severe heating, the research investigation entitled "Hygienic Collection and Preservation of Neera from Palmyra palm has been taken

Materials and Methods

Collection and preservation of neera

There were two experiments conducted. The experiment I (Hygienic collection of neera) was carried out in the farmer's field situated within a distance of 4-5 km radius surrounding the university, Bihar Agricultural University (BAU), Sabour. The experiment II (Preservation of neera) was conducted in the laboratory, department of Food Science and Postharvest Technology, Bihar Agricultural College (BAC), Sabour, Bhagalpur. For conducting experiment I, neera was tapped/collected from farmer's field in the early morning i.e. before sunrise. It was collected as per treatment details (Table-1).

Table 2. Parameters for selection of best method of neera collection

Parameters	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆
Alcohol percentage (% v/v)						
pH						
Titrateable acidity (%)						
Total Microbial Count (CFU/mL)						

The least fermented neera in terms of minimum % alcohol, minimum change in pH, and titrateable acidity (%) and least total microbial count (Cfu/mL) was rated as the best tapping treatment or method.

EXPERIMENT II: Preservation of neera

For conducting experiment II, the neera was collected by the best treatment method identified in experiment I. Treatment T₆ was identified as the best neera collection method where an specially designed box was used. As per treatment details of 2nd experiment, (Table-3), different preservation methods were used for preserving fresh neera. After treatments, neera was packed in 200 mL PET bottle and stored at 2-3°C and quality analysis parameters were recorded initially and after an interval of 15 days for a period of two months. The sensory quality attributes were also evaluated at the same interval. The evaluation of sensory attributes were based on nine-point hedonic scale (Annexure I).

Table 3. Experiment -II: Treatment Details (Preservation of Neera)

Treatments	Treatment details
T ₁	No Pasteurization (Control)
T ₂	Pasteurization at 60°C for 10 minutes
T ₃	Pasteurization at 65°C for 10 minutes
T ₄	Treatment with ultrasonic (30 kHz) for 10 minutes + Pasteurization at 60°C for 10 minutes
T ₅	Treatment with ultrasonic (30 kHz) for 10 minutes + Pasteurization at 65°C for 10 minutes
T ₆	Treatment with ultrasonic (35 kHz) for 15 minutes + Pasteurization at 65°C for 10 minutes
T ₇	Treatment with ultrasonic (35 kHz) for 20 minutes + Pasteurization at 65°C for 10 minutes

Analysis of fresh neera

Chemical parameters

The physico-chemical parameters were analysed as per standard procedure. Titrateable acidity of neera was estimated as per AOAC, [19]. Total soluble solids of neera was estimated by using hand refractometer (Erma hand refractometer with automatic temperature controller, ERB-32, S/N: 600/B19/36) and the results are expressed as degree brix (°B). The pH of the neera was determined with the help of digital pH meter. The pH meter was first standardized with buffers of pH 4.0 and 7.0 before pH measurements of the samples. The percent total and reducing sugars of neera were estimated by Lane and Eynon's method [19] by titrating a known volume of the samples against Fehling's solutions (Fehling A and Fehling B) using methylene blue as indicator. Total phenol content was estimated by using the Folin-Ciocalteu reagent [20].

Microbial analysis

For analyzing microbial load (yeast, mould and bacteria) in neera two different media, (Potato dextrose agar (PDA) and nutrient agar (NA) were used. Potato dextrose agar (PDA) media was used for counting fungal and yeast growth and nutrient agar (NA) was used for bacterial count. The media (PDA and NA) was prepared as described by Opara et al. [21] with required modifications. All laboratory grade chemicals were used for analysis.

Organoleptic evaluation

Organoleptic analysis was carried out by judges comprising of PG students, teaching and non-teaching staffs (six number of judges) as per methods described by Ranganna [22]. The evaluation was for taste and overall acceptability on 9-point Hedonic scale as mentioned in annexure II.

Statistical analysis

The data of 1st experiment (experiment I) was analysed by Randomized Block Design (RBD) and 2nd by Controlled Randomized Design (CRD), respectively.

The data were analyzed by analysis of variance (ANOVA) and means were compared by Duncan's Multiple Range Test (DMRT) using SPSS Version 26. Differences between the means at 5 % level were considered significant.

Results

Bio-chemical and microbial estimation of fresh neera

The bio-chemical characteristics of fresh neera as influenced by the tapping method is presented in Table 4. It is clear from the table that the tapping methods affected biochemical parameters. The lowest TSS was observed in control (T₁) (9.50±0.50°B) and highest in T₆ (12.33±0.29°B) followed by T₅ (12.00±1.80°B). The titrateable acidity (%) of control (T₁) was found significantly higher from rest of the treatments. The highest titrateable acidity (%) was observed in T₁ (1.05±0.18%) and the lowest in T₂ followed by T₄. However, except T₁, all other treatments were found non significant. The pH value of control (T₁) was also found significantly lower from T₅ and T₆. The minimum pH was observed in T₁ (3.94±0.52) and maximum in T₅ (5.02±0.41) followed by T₆ (4.67±0.41). Alcohol percentage was found significantly higher in control sample (5.17±0.58%) as compared to T₆ (3.93±0.35%). Percent reducing sugar was found lowest in control sample and significantly different from T₆ and T₇ treatments but non-significant with other treatments. Similarly, percent total sugar was also found significantly lower in control treatment as compared to rest of the treatments except T₂ which was found at par with T₂. The phenol (mg/100 mL) content was found non significant.

The microbial population in fresh neera is very much important because it is directly related to fermentation and spoilage. The maximum microbial load (bacteria, yeasts and moulds) was recorded in control sample (T₁) and bacterial population was high compared to yeast (figure 1). The best collection treatment was found to be T₆ (Sanitization with AFS* Solution (1500 ppm So₂ +0.2% C.A.) + Plastic Pot lime (0.5%) + protected by cushioning with thermocol+ 300 g Crushed ice) in terms of less microbial load, low acidity, low alcohol percentage and high pH as compared to other treatments.

The acceptability of neera depends upon the organoleptic parameters like taste, odour and over all acceptability (figure 2). It is evident from the figure that there is a significant difference in all three organoleptic parameters (taste, odour and over all acceptability) of fresh neera on 9-point hedonic scale. The highest score (7.33) equivalent to like moderately was observed in T₄ and T₆ and minimum in control sample (3.67±0.58) which was equivalent to dislike moderately. The lowest score was registered in control sample (T₁) and highest in T₆. Other treatments, like T₂ to T₅ were found significantly superior than control but at par to each other. On the other hand, anti fermenting solution that was applied in T₆ (Sanitization with AFS* Solution (1500 ppm So₂ + 0.2% C.A.) + Plastic Pot lime (0.5%) + protected by cushioning with thermocol+ 300 g Crushed ice) was found most effective and so the OAA taste was highest in T₆ as compared to others. Hence hygienic collection of the neera preserves the quality for comparatively longer duration [23].

Table 4. Bio-chemical characteristics of fresh neera as influenced by different tapping methods

Treatments (T)	TSS (°B)	Acidity (%)	pH	Alcohol (%)	Reducing Sugar (%)	Total Sugar (%)	Phenol (mg/100 mL)
T ₁	9.50±0.50 ^b	1.05±0.18 ^a	3.94±0.52 ^c	5.17±0.58 ^a	3.68±0.77 ^b	6.70±0.82 ^c	342.43±16.98 ^a
T ₂	10.25±0.75 ^b	0.54±0.10 ^b	4.34±0.17 ^{bc}	4.45±0.78 ^{ab}	3.76±0.70 ^b	7.42±0.66 ^{bc}	320.03±62.57 ^a
T ₃	10.32±0.12 ^b	0.64±0.23 ^b	4.25±0.08 ^{bc}	4.35±0.17 ^{ab}	3.87±0.36 ^b	8.73±0.34 ^a	310.77±175.52 ^a
T ₄	10.67±0.58 ^{ab}	0.60±0.05 ^b	4.44±0.26 ^{bc}	4.34±0.11 ^{ab}	4.40±0.15 ^{ab}	8.75±0.30 ^a	324.64±78.83 ^a
T ₅	12.00±1.80 ^a	0.65±0.10 ^b	5.02±0.41 ^a	4.33±0.13 ^{ab}	4.32±0.50 ^{ab}	8.63±0.45 ^a	323.93±134.71 ^a
T ₆	12.33±0.29 ^a	0.65±0.10 ^b	4.67±0.41 ^{ab}	3.93±0.35 ^b	4.90±0.12 ^a	8.13±0.31 ^a	307.79±81.99 ^a
T ₇	12.29±0.21 ^a	0.64±0.10 ^b	4.66±0.41 ^{ab}	3.83±0.35 ^b	4.81±0.12 ^a	8.32±0.31 ^a	306.79±81.99 ^a
SEm (±)	0.509	0.071	0.160	0.250	0.284	0.322	61.431
CD (P=0.05)	1.604	0.224	0.504	NS	1.021	1.015	NS

Table 5. Microbial population of fresh neera.

Treatment	Yeast population	Mould population	Bacterial Population
T ₁	2x 10 ⁶	1 x 10 ⁴	3 x 10 ⁶
T ₂	2x 10 ⁵	1 x 10 ³	3 x 10 ⁴
T ₃	2x 10 ⁴	1 x 10 ³	3 x 10 ³
T ₄	2x 10 ³	1 x 10 ²	3 x 10 ⁴
T ₅	2x 10 ³	1 x 10 ³	3 x 10 ⁴
T ₆	2x 10 ²	1 x 10 ³	3 x 10 ⁴
T ₇	2x 10 ²	1 x 10 ²	3 x 10 ³




Fig. Microbial population of fresh neera. Population in Log10 transformed value.

Table 6. Organoleptic attributes of fresh neera

Treatments (T)	Taste	Odour	Overall Acceptability
T ₁	3.67±0.58 ^c	5.00±1.00 ^c	4.67±0.58 ^c
T ₂	6.00±1.00 ^b	6.00±1.00 ^{bc}	6.00±1.00 ^b
T ₃	6.33±0.58 ^{ab}	6.33±0.58 ^{ab}	6.33±0.58 ^b
T ₄	7.33±0.58 ^a	7.33±0.58 ^a	6.67±0.58 ^b
T ₅	5.33±0.58 ^b	5.33±0.58 ^{bc}	5.67±0.58 ^{bc}
T ₆	7.33±0.58 ^a	6.00±1.00 ^{bc}	8.00±0.00 ^a
T ₇	7.31±0.58 ^a	6.00±1.00 ^{bc}	8.00±0.00 ^a
SEm (±)	0.380	0.350	0.360
CD (P=0.05)	1.197	1.103	1.134




Figure 1. Organoleptic attributes of fresh neera

Effect of preservation methods on biochemical parameters during storage at low temperature (2-3°C)

Under 2nd experiment, the treated neera was packed and stored at low temperature (2-3°C) for quality evaluation at a fixed interval of fifteen days for a total period of 60 days (2-months). During this period, biochemical and sensory observations were recorded. The following parameters were studied during storage.

Titrateable Acidity (%) and pH

In general, a successive decreasing pattern in titrateable acidity (%) was observed in all treatments from the start of the storage period till the end i.e. up to 60 days (table 7) except few treatments (T₁, T₂, T₃ and T₄) that spoiled before 15, 30 and 45 days, respectively. This decrease in acidity might be the result of acid-mediated hydrolysis of polysaccharides and non-reducing sugars to their simpler components where acid was utilized for converting them to hexose sugars or complexes in the presence of metal ions [24].

The pH value was found decreased in T₁, T₂, T₃ and T₄ but an increasing trend was observed in T₅, T₆ and T₇ treatments. The decrease in T₁, T₂, T₃ and T₄ might be due to initial lactic acid production by the process of natural fermentation.

Total Soluble Solids (°B), Reducing and Total Sugar (%)

The value of total soluble solids (TSS) is presented in figure 2 and that of sugars (reducing and total sugar) are presented in table 8. It is clear from the figure and table that four treatments, (T₁, T₂, T₃ and T₄) were rejected after noticing spoilage at different storage intervals. T₁ (control) was rejected after 2 days storage only whereas, T₂ after 15 days, T₃ after 30 days and T₄ after 45 days, respectively. Only three treatments namely T₅, T₆ and T₇, successfully stored for 60 days. The TSS and both reducing and total sugars of neera changed significantly during storage of 60 days. The initial value of TSS (°B) of T₅, T₆ and T₇ were found 12.00±1.80, 12.33±0.29 and 12.37±0.30°B, respectively which was found to be increased during 60 days of storage. The initial value of reducing sugar (%) of T₅, T₆ and T₇, were recorded 4.32±0.50%, 4.90±0.12% and 4.87±0.16% which was found to increase during 60 days of storage up to 5.09±0.13%, 5.25±0.05 and 5.17±0.08, respectively.

Table 7. Titratable acidity (%) and pH of treated neera during 60 days storage at low temperature (2-3oC)

Days Treatments	Titratable acidity (%)						pH					
	0-day	15-days	30-days	45-days	60-days	Mean	0-day	15-days	30-days	45-days	60-days	Mean
T ₁	1.05±0.18	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.21±0.44^d	3.94±0.52	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.79±1.64^f
T ₂	0.54±0.10	0.82±0.06	0.00±0.00	0.00±0.00	0.00±0.00	0.27±0.36^d	4.34±0.17	3.88±0.13	0.00±0.00	0.00±0.00	0.00±0.00	1.64±2.09^e
T ₃	0.64±0.23	0.55±0.15	0.93±0.28	0.00±0.00	0.00±0.00	0.42±0.41^c	4.25±0.08	4.27±0.10	3.76±0.16	0.00±0.00	0.00±0.00	2.46±2.09^d
T ₄	0.60±0.05	0.58±0.06	0.55±0.10	0.75±0.10	0.00±0.00	0.50±0.27^b	4.44±0.26	4.55±0.19	4.69±0.13	3.71±0.53	0.00±0.00	3.48±1.85^c
T ₅	0.65±0.10	0.58±0.06	0.53±0.04	0.48±0.08	0.47±0.08	0.54±0.09^{ab}	5.02±0.41	5.09±0.30	5.25±0.52	5.37±0.36	5.46±0.28	5.24±0.37^a
T ₆	0.65±0.10	0.57±0.16	0.53±0.10	0.50±0.05	0.42±0.03	0.53±0.12^{ab}	4.67±0.41	4.78±0.49	5.00±0.12	5.11±0.10	5.22±0.22	4.96±0.34^b
T ₇	0.68±0.06	0.63±0.10	0.60±0.04	0.56±0.02	0.48±0.06	0.59±0.09^a	4.93±0.54	5.12±0.49	5.13±0.19	5.24±0.31	5.45±0.2	5.18±0.36^a
Mean	0.69±0.19^a	0.53±0.25^b	0.45±0.34^c	0.33±0.31^d	0.20±0.23^e		4.51±0.48^a	3.95±1.73^b	3.41±2.26^c	2.78±2.53^d	2.3±2.73^e	
	Days (D)		Treatment (T)		D × T		Days (D)		Treatment (T)		D × T	
SEm (±)	0.021		0.025		0.055		0.060		0.071		0.160	
CD (P=0.05)	0.059		0.071		0.155		0.169		0.200		0.451	




Figure 2. Total Soluble Solids (TSS) (°B) of treated neera during 60 days storage at low temperature (2-3oC)




Figure 3. Total Phenol (mg/100 mL) of treated neera during 60 days storage at low temperature (2-3oC)

Total Phenol (mg/100 mL) and Organoleptic properties of neera

The value of total phenol was found in decreasing order till the end of storage (Figure 3). The decrease was observed in all treatments. The initial value of Total phenol (mg/100 mL) of the best treatments (T₅, T₆ and T₇) were recorded as 323.92±134.73, 307.80±81.98 and 323.73±55.96 (mg/100 mL) which has been reduced to 223.82±50.10, 223.91±51.42 and 225.63±45.7 (mg/100 mL), respectively, at the end of 60 days storage.

Table 8. Reducing and Total Sugars (%) of treated neera during 60 days storage at low temperature (2-3°C)

Days Treatments	Reducing Sugars (%)						Total Sugars (%)					
	0-day	15-days	30-days	45-days	60-days	Mean	0-day	15-days	30-days	45-days	60-days	Mean
T ₁	3.68±0.77	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	0.74±1.55 ^f	6.70±0.82	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	1.34±2.79 ^f
T ₂	3.76±0.70	3.94±0.58	0.00±0.00	0.00±0.00	0.00±0.00	1.54±1.98 ^e	7.42±0.66	7.52±0.63	0.00±0.00	0.00±0.00	0.00±0.00	2.99±3.80 ^e
T ₃	3.87±0.36	4.03±0.53	4.33±0.15	0.00±0.00	0.00±0.00	2.44±2.09 ^d	8.73±0.34	9.01±0.18	9.13±0.25	0.00±0.00	0.00±0.00	5.38±4.55 ^d
T ₄	4.40±0.15	4.48±0.14	4.59±0.05	4.75±0.10	0.00±0.00	3.64±1.89 ^c	8.75±0.30	8.85±0.29	9.03±0.24	9.03±0.24	0.00±0.00	7.13±3.70 ^c
T ₅	4.32±0.50	4.67±0.13	4.74±0.07	4.83±0.08	5.09±0.13	4.73±0.33 ^b	8.63±0.45	8.77±0.39	8.89±0.33	8.89±0.33	8.97±0.33	8.83±0.33 ^a
T ₆	4.90±0.12	4.90±0.12	5.01±0.12	5.10±0.17	5.25±0.05	5.03±0.17 ^a	8.13±0.31	8.23±0.38	8.45±0.51	8.45±0.51	8.68±0.52	8.39±0.43 ^b
T ₇	4.87±0.16	4.84±0.06	4.97±0.25	4.93±0.24	5.17±0.08	4.96±0.19 ^a	8.06±0.3	8.12±0.33	8.27±0.52	8.23±0.61	8.57±0.5	8.25±0.43 ^b
Mean	4.26±0.62 ^a	3.84±1.66 ^b	3.38±2.20 ^c	2.8±2.49 ^d	2.22±2.62 ^e		8.06±0.84 ^a	7.21±3.07 ^b	6.25±4.07 ^c	4.94±4.40 ^d	3.75±4.44 ^e	
	Days (D)		Treatment (T)		D × T		Days (D)		Treatment (T)		D × T	
SEm (±)	0.057		0.068		0.152		0.081		0.095		0.213	
CD (P=0.05)	0.161		0.192		0.429		0.228		0.268		0.601	

Effect of preservation methods on sensory qualities of neera during storage at low temperature (2-3°C)

The initial taste and overall acceptability (OAA) score of treatments, T₁ to T₄ was registered as lower values (6-7 approx) on hedonic scale confirming that the fermentation started in these treatments and spoiled within few days (table 9). On the other hand, higher scores have been registered in T₅ (6.67±1.15), T₆ (7.33±0.58), T₇ (8.00±0.00) for taste, and T₅ (5.677.00±1.00), T₆ (8.00±0.00), T₇ (8.33±0.58) for OAA, respectively. The final taste score of T₅, T₆, and T₇ were found to be 6.00±1.00, 6.67±0.58 and 7.33±0.58 for taste and 6.00±0.00, 6.67±0.58 and 7.00±0.00 for OAA, respectively. This indicates that both taste and OAA score decreased with the passage of time. Fermentation started immediately in the control sample (T₁) followed by T₂, T₃ and T₄ and hence scored lower values on hedonic scale. On the other hand, higher scores have been registered in T₅, T₆ and T₇ for both taste and overall acceptability (OAA). The final taste score of T₅, T₆, and T₇ was acceptable even after 60 days of storage but loss in both taste and OAA was observed.

Table 9. Taste and Overall acceptability (OAA) of treated neera during 60 days storage at low temperature (2-3°C)

Days Treatment	Taste						Overall acceptability					
	0-day	15-days	30-days	45-days	60-days	Mean	0-day	15-days	30-days	45-days	60-days	Mean
T ₁	6.33±1.15	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	1.27±2.66 ^f	6.00±1.00	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00	1.20±2.51 ^f
T ₂	6.67±0.58	4.33±0.58	0.00±0.00	0.00±0.00	0.00±0.00	2.20±2.91 ^e	6.00±1.00	4.67±0.58	0.00±0.00	0.00±0.00	0.00±0.00	2.13±2.77 ^e
T ₃	7.33±0.58	6.67±0.58	3.67±0.58	0.00±0.00	0.00±0.00	3.53±3.27 ^d	6.33±0.58	6.33±0.58	4.33±0.58	0.00±0.00	0.00±0.00	3.40±3.00 ^d
T ₄	7.33±0.58	7.00±0.00	6.67±0.58	4.33±0.58	0.00±0.00	5.07±2.87 ^c	6.67±0.58	6.67±0.58	6.33±0.58	5.00±1.00	0.00±0.00	4.93±2.69 ^c
T ₅	7.33±0.58	6.67±0.58	6.67±0.58	6.00±0.00	6.00±1.00	6.53±0.74 ^b	7.00±1.00	7.00±0.00	6.67±0.58	6.67±0.58	6.00±0.00	6.67±0.62 ^b
T ₆	7.67±0.58	7.33±0.58	7.33±0.58	7.33±0.58	7.33±0.58	7.40±0.51 ^a	8.00±0.00	8.00±0.00	7.67±0.58	7.00±0.00	6.67±0.58	7.47±0.64 ^a
T ₇	8.00±0.00	7.67±0.58	7.67±0.58	7.67±0.58	7.67±0.58	7.73±0.46 ^a	8.33±0.58	7.67±0.58	7.33±0.58	7.33±0.58	7.00±0.00	7.53±0.64 ^a
Mean	7.24±0.77 ^a	5.67±2.61 ^b	4.57±3.23 ^c	3.62±3.38 ^d	3.00±3.61 ^e		6.90±1.09 ^a	5.76±2.64 ^b	4.62±3.19 ^c	3.71±3.39 ^d	2.81±3.34 ^e	
	Days (D)		Treatment (T)		D × T		Days (D)		Treatment (T)		D × T	
SEm (±)	0.111		0.131		0.293		0.110		0.130		0.292	
CD (P=0.05)	0.313		0.369		0.826		0.310		0.367		0.824	

Discussion**Effect of hygienic collection on biochemical, microbial and sensory qualities of fresh neera**

It is clear from the table 4 and 5 that the hygienic collection of neera controls changes in biochemical parameters and microbial load of fresh neera, respectively. Biochemical parameters like total soluble solids, percent acidity, pH, percent alcohol, phenol, total and reducing sugars content were found different in different treatments. The titratable acidity (%) of control (T₁) was found significantly different from rest of the treatments. The minimum pH was observed in T₁ (3.94±0.52) and maximum in T₅ (5.02±0.41) followed by T₆ (4.67±0.41). High acid and low pH observed in control sample might be due to lactic acid fermentation [25].

Lasekan et al. also observed higher percentage of total acidity and low pH in ten palm sap samples collected from Songkhla province of Southern Thailand. The lowest total soluble solids (TSS) was observed in control (T₁) (9.50±0.50°B) and highest in T₆ (12.33±0.29°B) followed by T₅ (12.00±1.80°B). Alcohol percentage and microbial population were found significantly higher in control sample as compared to T₆. This might be due to the unhygienic collection of neera that might have increased microbial contamination and increased fermentation rate in control sample. The increase in fermentation rate converted more sugar content into alcohol, resulted decrease in TSS content and an increase in alcohol content. Kurniawan et al., [26] also reported that the degree of fermentation of the palmyrah neera depends on the practice of neera collection and the environmental conditions.

Hygiene collection of the neera samples preserves it in fresh form for a comparatively longer duration [23]. The maximum value of microbial population in control sample (T_1) be due to unhygienic collection of neera. However, bacterial population was found more than yeast and moulds. This might be due to more bacterial contamination than yeast and moulds. According to Somashekaraiah et al. [11], neera has been reported to have about 75 distinct Lactic acid bacteria (LAB) strains.

The value of organoleptic attributes like taste, odour and over all acceptability (OAA) were found significantly different on 9-point hedonic scale (table 6). The lowest score registered in control sample (T_1) and highest in T_6 . The reason might be the microbial contamination in control sample where anti fermenting solution (AFS) was not applied. On the other hand, the most effective anti fermenting solution was applied in T_6 (Sanitization with AFS* Solution (1500 ppm SO_2 + 0.2% C.A.) + Plastic Pot lime (0.5%) + protected by cushioning with thermocol + 300 g Crushed ice). Hence hygiene collection of the neera preserves it in fresh form for a comparatively longer duration [23]. Kurniawan et al., [26] also reported that the degree of fermentation of the palmyraneera depends on the methods of neera collection and the environmental conditions.

Effect of preservation methods on shelf life of neera at low temperature (2-3°C)

The aim of 2nd experiment was to identify the best treatment that can prevent spoilage up to 60 days. The control sample (T_1) spoiled within 3 days. Treatments like T_2 , T_3 , and T_4 spoiled with 15, 30 and 45 days, respectively. The thermosensation treatments, T_5 , T_6 and T_7 , only completed 60 days. Therefore, it was concluded that these three treatments (ultrasonic (30 kHz) for 10 minutes + Pasteurization at 65°C for 10 minutes (T_5), ultrasonic (35 kHz) for 15 minutes + Pasteurization at 65°C for 15 minutes (T_6), ultrasonic (35 Hz) for 15 minutes + Pasteurization at 65°C for 20 minutes (T_7)) were effective treatments in preventing fermentation and spoilage of neera. It has been reported that thermosonication has higher efficiency for killing of microbes and deactivation of enzymes at relatively low temperatures [27]. Kapturowska et al. [18] also reported that thermosonication is very effective against yeast.

Effect of preservation methods on biochemical parameters during storage at low temperature (2-3°C)

After 2nd experiment, the samples were stored at low temperatures and again biochemical changes and sensory qualities of neera such as titratable acidity (%), pH, percent reducing and total sugars, total soluble solids (°B), total phenol (mg/100 mL), taste and overall acceptability were analyzed during the storage period of 60 days. In general, a successive decreasing pattern in titratable acidity (%) and percent total phenol were observed.

This decrease in acidity might be the result of acid-mediated hydrolysis of polysaccharides and non-reducing sugars to their simpler components where acid was utilized for converting them to hexose sugars or complexes in the presence of metal ions [24]. Similarly, the decrease in polyphenol content might be either due to various polyphenol mediated reactions during storage or polyphenol compounds might have contributed to haze formation by interacting with proteins through polymerization or oxidation leading to the formation of high-molecular-weight polymeric complexes [28]. At the same time, an increasing pattern was observed in pH, total soluble solids, percent reducing and total sugars in all treatments from the

starting of storage period till the end i.e. up to 60 days. Some of the treatments like T_1 , T_2 , T_3 and T_4 not followed the general trend and were spoiled before 15, 30 and 45 days, respectively. In these treatments, an increase in percent acidity and decrease in pH were recorded.

The decrease in acidity and a corresponding increase in pH might be due to fermentation of neera before spoilage. Leena et al. [29] also reported an increase in percent titratable acidity and decrease in pH of palmyrah palm neera due to microbial fermentation of the carbohydrates present in the palmyrah neera. Prashanth and Patil [30] also reported a increase in acidity and a decrease in pH of coconut sap during storage.

The increase in TSS might be due to hydrolysis of polysaccharides and pectic substances into simpler soluble saccharides which might have contributed to an increase in TSS during storage [24]. A nearly similar result was also observed by Nath et al. [31] in ginger- kinnow squash and in Kinnow mandarin juice blends [32]

The increase in reducing sugar (%) might be due to hydrolysis of polysaccharides mainly starch and pectic substances into simpler soluble saccharides which contributed an increase in reducing and total sugar (%) during storage [24]. Naknean et al. [33] reported an increase in reducing sugar due to inversion of sucrose into glucose and fructose. Rao et al. [34] also reported an increase in %reducing and total sugar in palmyra neera during four months of storage and then declined. Naknean et al [35] also reported an increase in reduced sugar of palmyra palm sap during storage after treatment with clarifying agents. Yousaf et al. [36] and Singh et al. [37] also reported increase in reducing sugar during storage of banana juice and processed sugar can juice, respectively.

Effect of preservation methods on sensory qualities of neera during storage at low temperature (2-3°C)

Fermentation started immediately in the control sample (T_1) followed by T_2 , T_3 and T_4 and hence scored lower values on hedonic scale. On the other hand, higher scores have been registered in T_5 , T_6 and T_7 for both taste and overall acceptability (OAA). The final taste score of T_5 , T_6 and T_7 was acceptable even after 60 days of storage but loss in both taste and OAA was observed. The loss of taste and over all acceptability might be due to undesirable biochemical changes [38] or the degradation of ascorbic acid and furfural production [39]. A similar trend is also reported by Rao et al. [34] during six weeks of storage at 2-4°C. The results are in accordance with [40] in coconut water and [41] in palmyra neera.

Summary and Conclusion

For hygienic tapping of neera from palmyrah palm (experiment I), the treatment T_6 (Sanitization with AFS* Solution (1500 ppm SO_2 + 0.2% C.A.) + Plastic Pot lime (0.5%) + protected by cushioning with thermocol + 300 g crushed ice) was selected as the best treatment and carry forwarded for conducting experiment II. Among different treatments of experiment II, treatments T_6 and T_7 , (ultrasonic (35 kHz) for 15 minutes + Pasteurization at 65°C for 10 minutes and ultrasonic (35 kHz) for 20 minutes + Pasteurization at 65°C for 10 minutes Pasteurization, respectively, were found the best treatments for preservation of neera at low temperature on the basis of no fermentation, longer shelf life, least changes in biochemical and sensorial qualities. Prashanth and Patil (2020) also reported that pasteurization increases the shelf life of coconut sap.

Hence treatment T₆ of experiment I and treatments T₆ and T₇ of experiment II may be commercially exploited for the safe and hygienic collection and preservation of neera from palmyrah palm.

Acknowledgment

We acknowledge the help extended by other departments of Bihar Agricultural University (BAU), Sabour particularly the Department of Plant Pathology, BAU, Sabour. We wish to express our sincere thanks to all helped us directly or indirectly in the collection of neera in the early morning, transportation, and analysis.

Disclosure statement

There is no conflict of interest among the authors.

Funding

This research work was conducted as part of a postgraduate (M.Sc.) thesis funded by Bihar Agricultural University (BAU), Sabour, Bhagalpur, Bihar, India.

References

1. Theivendirarajah K (2008) Palmyrah Palm – A Monograph, Department of Botany, University of Jaffna, Sri Lanka.
2. Rao M C, Swami D V, Pandit A, Nanda S P and Rao B B (2021) Scope, Nutritional Importance and Value Addition in Palmyrah (*Borassus flabellifer* L.): An Under Exploited Crop. Bioactive Compounds: Biosynthesis, Characterization and Applications, 29:207. <http://dx.doi.org/10.5772/intechopen.91131>.
3. Veilmuthu P (2020) Palmyra – nature's perennial gift in the face of climate crisis. Available: <http://climate.southasia.org/palmyra-natures-perennial-gift-in-the-face-ofclimate-crisis/> (Accessed on 04.07.2020)
4. Upadhyaya A and Sonawane S K (2023) Palmyrah palm and its products (Neera, Jaggery and Candy) - A Review on chemistry and technology. Applied Food Research 3 (1) 100256. <https://doi.org/10.1016/j.afres.2022.100256>
5. Vengaiya P C V, Murthy G N, Sattiraju M and Maheswarappa H P (2017) Value added food products from Palmyrah palm (*Borassus flabellifer* L). Journal of Nutrition and Health Sciences, 4(1): 105. doi: 10.15744/2393-9060.4.105
6. Gummadi V P, Battu G R, Keerthana, Diyya M S and Manda K (2016) A review on palmyrah palm (*Borassus flabellifer*). International Journal of Current Pharmaceutical Research, 8 (2):1720. https://www.researchgate.net/publication/316887881_A_review_on_palmyra_palm_Borassus_flabellifer
7. Kalimuthu K (2019) Neera and its importance-a review.10.24327/ijrsr.2019.1002.3030.
8. Hebbar K B, Arivalagan M, Manikantan M R, Mathew A C, Thamban C, Thomas G V and Chowdappa P (2015) Coconut inflorescence sap and its value addition as sugar – collection techniques, yield, properties and market perspective. Current Science, 109(8), 1411–1417. <http://www.jstor.org/stable/24905994>
9. Saranya P (2016) Characterization of palmyra fruit *Borassus flabellifer* linn pulp and development of ready to serve beverage from palmyra fruit pulp. Doctoral dissertation, Periyar University, Salem, India; Available: <https://shodhganga.inflibnet.ac.in/handle/10603/235390>
10. Kurniawan T, Jayanudin Kustiningsih I and Firdaus M A (2018) Palm sap sources, characteristics, and utilization in Indonesia. Journal of Food and Nutrition Research 6(9): 590-596. <https://doi.org/10.12691/jfmr-6-9-8>
11. Somashekaraiah R, Shruthi B, Deepthi B V and Sreenivasa M Y (2019) Probiotic properties of lactic acid bacteria isolated from Neera: A naturally fermenting coconut palm nectar. Frontiers in Microbiology, 10. <https://doi.org/10.3389/fmicb.2019.01382>
12. Alonso S, Carmen C. M, Berdasco M, de la Banda I G, Moreno-Ventas X and de Rojas A H (2019). Isolation and partial characterization of lactic acid bacteria from the gut microbiota of marine fishes for potential application as probiotics in aquaculture. Probiotics and antimicrobial proteins, 11, 569-579. <https://doi.org/10.1007/s12602-018-9439-2>
13. Kurian A and Peter K V (2007) Commercial Crops Technology. Horticulture Science Series: 8. New India Publishing.
14. Tulashie SK, Appiah A P, Torku G D, Darko A Y and Wiredu A (2017) Determination of methanol and ethanol concentrations in local and foreign alcoholic drinks and food products (Banku, Gakenkey, Fantekenkey and Hausa koko) in Ghana. Food Contamination 4(1): 1-5. <https://doi.org/10.1186/s40550-017-0059-5>
15. Arias C, Murray B (2012) The rise of the Enterococcus: beyond vancomycin resistance. Nat Rev Microbiology 10, 266–278. <https://doi.org/10.1038/nrmicro2761>
16. Ho C W, Wan A M W, Maskat Y M and Osman H (2007) Changes in volatile compounds of palm sap (*Arenga pinnata*) during the heating process for production of palm sugar. Food Chemistry, 102: 1156–1162. <https://doi.org/10.1016/j.foodchem.2006.07.004>
17. Ercan S Ş and Soysal Ç (2013) Use of ultrasound in food preservation, Natural Science, 05 (8A2): 5–13. doi:10.4236/ns.2013.58A2002.
18. Kapturowska A, Stolarzewicz I and Chmielewska I (2011) Ultrasounds—A tool to inactivate yeast and to extract intracellular protein. Zywnosc-Nauka Technologia Jakosc, 18, 160-171.
19. AOAC (2000) Official Methods of Analysis. Association of Official Analytical Chemists, Washington, DC, USA.
20. Sadasivam M S and Manickam A (2005) Biochemical Methods. Newage International Publishers. Second Edition. 193p.

21. Opara C C, Ajoku G and Madumelu N O (2014) Palm wine mixed culture fermentation kinetics. Greener Journal of Physical Sciences 3: 28-37. SOI: <http://s-o-i.org/1.15/ijarbs-2-12-6>
22. Ranganna S (2014) Handbook of Analysis and Quality Control for Fruits and Vegetable Products. 12nd edn. Tata Mc Graw Hill Publishing Co. Ltd., New Delhi, India (21st reprint 2014)
23. Khandekar R G, Maheswarappa H P and Sanapa P B (2018) Inter cropping of spices in coconut garden-A success story from Maharashtra. Indian Coconut Journal 61(1): 4-5. <http://coconutboard.in>
24. Bohra P, Sreenivas N K and Sreeramu B S (2012) Development of a cost-effective, palatable and shelf-stable blended beverage of pummelo (*Citrus grandis* Linn.) Fruits 67(4):249-256. DOI: <https://doi.org/10.1051/fruits/2012018>
25. Lasekan O, Buettner A and Christlbauer M (2007) Investigation of important odorants of palm wine (*Elaeis guineensis*). Food Chemistry 105(1): 15-23 <https://doi.org/10.1016/j.foodchem.2006.12.052>.
26. Kurniawan T, Kustiningsih I and Firdaus M A (2020), Palm sap sources, characteristics, and utilization in Indonesia. Journal of Food and Nutrition Research 6 (9): 590-596. <http://pubs.sciepub.com/jfmr/6/9/8>. DOI:10.12691/jfmr-6-9-8.
27. Ercan S Ş and Soysal Ç (2013) Use of ultrasound in food preservation, Natural Science, 05 (8A2): 5-13. doi:10.4236/ns.2013.58A2002.
28. Lee M Y, Lee M K and Park I (2007) Inhibitory effect of onion extract on polyphenol oxidase and enzymatic browning of taro (*Colocasia antiquorum* var. esculenta). Food Chemistry 105(2): 528-532. <https://doi.org/10.1016/j.foodchem.2007.04.010>
29. Leena M M, Yoha K S, Moses J A and Anandharamakrishnan C (2021) Electrospun nanofibrous membrane for filtration of coconut neera. Nanotechnology for Environmental Engineering 6(2): 1-10. <https://doi.org/10.1007/s41204-021-00116-1>
30. Prashanth B K and Patil V S (2020) Effect of thermal pasteurization on shelf life & quality of Kalparasa (Coconut sap). Journal of Farm Sciences, 33(1), 131-134. <https://d1wqtxts1xzle7.cloudfront.net/65592269/2018380>
31. Nath A, Yadav D S, Sarma P and Dey B (2005) Standardization of ginger - kinnow squash and its storage. Journal of Food Science and Technology 42(6): 520-522. <https://www.researchgate.net/publication/296714080>
32. Bhardwaj R and Nandal U (2014) Effect of storage temperature on physico-chemical and sensory evaluation of kinnow mandarin juice blends. Journal of Food Processing Technology 5:8, 1-4. <http://dx.doi.org/10.4172/2157-7110.1000361>
33. Naknean P, Juntorn N and Yimyuan T (2014) Influence of clarifying agents on the quality of pasteurised palmyra palm sap (*Borassus flabellifer* Linn.) International Journal of Food Science and Technology, 49: 1175-1183. <https://doi.org/10.1111/ijfs.12414>
34. Rao M C S, Swami D V, Ashok P, Suneetha D R S, Sujatha R V and Sekhar V (2022) Influence of different storage temperatures and packing material in extending shelf life and quality attributes of palmyrah (*Borassus flabellifer* L.) neera. Journal of Plantation Crops, 50(2): 85-93. doi: 10.25081/jpc.2022.v50.i2.7982.
35. Naknean P, Jutasukosol K and Mankit T (2015) Utilization of chitosan as an antimicrobial agent for pasteurized palm sap (*Borassus flabellifer* Linn.) during storage. Journal of Food Science and Technology 52 , 731-741. <https://doi.org/10.1007/s13197-013-1104-x>
36. Yousaf M S, Yusaf S, Myba M and Abd-Aziz S (2010) Storage stability of clarified banana juice fortified with inulin and oligofructose. Journal of Food Processing and Preservation, 34 (2): 599-610. URL: <http://www.blackwell-synergy.com/loi/jfpp>
37. Singh I, Solomon S, Shirivastava A H, Singw R K and Singw J (2006) Post-harvest quality deterioration of cane juice: physiobiochemical indicators. Sugar Technology, 8, 128-131. <http://www.iapsit.org> DOI: 10.1007/BF02943646
38. Dewivedi S K, Pathak S and Mishra V (2011) Formulation and evaluation of RTS beverage from black mulberry. International Symposium on Minor Fruits and Medicinal Plants for Health and Ecological Security (ISMF & MP) Kalyani, West Bengal, 259-263.
39. Kausar H, Saeed S, Ahmad M M and Salam A (2012) Studies on the development and storage of cucumber-melon functional drink. Journal of Agricultural Research, 50(2):239-248. https://www.researchgate.net/profile/Muhammad-Ahmad-367/publication/329239908_Development_and_storage_stability_of_cucumbermelon_drink
40. Gobin A, Falade K O and Akingbala J O (2009) Effect of packaging on physical, chemical and sensory attributes of coconut water during storage. Journal on Food Agriculture and Environment 7(1): 62-65. URL: <http://www.isfae.org/scientificjournal/2009/issue1/pdf/food>
41. Anonymous (2013) Improving the shelf life of inflorescence sap. AICRP (Palms). Annual Report 2012-13. ICAR-All India Co-ordinated Research Project on Palms. ICARCPRI, Kasaragod. 84p.